

(2) *Swine*. A tolerance is established for parent doramectin (marker residue) in liver (target tissue) of 160 parts per billion.

[63 FR 68184, Dec. 10, 1998]

§ 556.227 Eprinomectin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of eprinomectin is 10 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle*. Tolerances are established for residues of eprinomectin B1a (marker residue) in milk of 12 parts per billion, in liver (target tissue) of 4.8 parts per million, and in muscle of 100 parts per billion.

(2) [Reserved]

[63 FR 59715, Nov. 5, 1998]

§ 556.228 Enrofloxacin.

The acceptable daily intake for enrofloxacin is 3 micrograms per kilogram of body weight per day.

(a) *Chickens and turkeys*. A tolerance of 0.3 part per million is established for residues of enrofloxacin (marker residue) in muscle (target tissue) of chickens and turkeys.

(b) *Cattle*. A tolerance of 0.1 part per million for desethyle ciprofloxacin (marker residue) has been established in liver (target tissue) of cattle.

[61 FR 56893, Nov. 5, 1996, as amended at 63 FR 49003, Sept. 14, 1998]

§ 556.230 Erythromycin.

Tolerances for residues of erythromycin in food are established as follows:

(a) 0.1 part per million in uncooked edible tissues of beef cattle and swine.

(b) Zero in milk.

(c) 0.025 part per million in uncooked eggs.

(d) 0.125 part per million (negligible residue) in uncooked edible tissues of chickens and turkeys.

[40 FR 13942, Mar. 27, 1975, as amended at 58 FR 43795, Aug. 18, 1993]

§ 556.240 Estradiol and related esters.

No residues of estradiol, resulting from the use of estradiol or any of the related esters, are permitted in excess of the following increments above the

concentrations of estradiol naturally present in untreated animals:

(a) In uncooked edible tissues of heifers, steers, and calves:

(1) 120 parts per trillion for muscle.

(2) 480 parts per trillion for fat.

(3) 360 parts per trillion for kidney.

(4) 240 parts per trillion for liver.

(b) In uncooked edible tissues of lambs:

(1) 120 parts per trillion for muscle.

(2) 600 parts per trillion for fat, kidney, and liver.

[49 FR 13873, Apr. 9, 1984, as amended at 56 FR 67175, Dec. 30, 1991]

§ 556.260 Ethopabate.

Tolerance for residues of ethopabate converted to metaphenetidine are established in the edible tissues of chickens as follows:

(a) 1.5 parts per million in uncooked liver and kidney.

(b) 0.5 part per million in uncooked muscle.

§ 556.270 Ethylenediamine.

A tolerance of zero is established for residues of ethylenediamine in milk.

§ 556.273 Famphur.

Tolerances are established for residues of famphur including its oxygen analog in or on meat, fat, or meat by-products of cattle at 0.1 part per million.

[62 FR 55161, Oct. 23, 1997]

§ 556.275 Fenbendazole.

(a) *Cattle and goats*. A tolerance¹ of 0.8 part per million is established for parent fenbendazole (the marker residue) in the liver of cattle and goats.

(b) *Swine*. A tolerance¹ for marker residues of fenbendazole in swine is not needed.

(c) *Cattle milk*. A safe concentration of 1.67 parts per million is established

¹As used in this section: "tolerance" refers to a concentration of a marker residue in the target tissue selected to monitor for total residues of the drug in the target animal.

for total fenbendazole residues. A tolerance of 0.6 part per million is established based on the fenbendazole sulf-oxide metabolite (marker residue).

[59 FR 26943, May 25, 1994, as amended at 61 FR 29478, June 11, 1996]

§ 556.277 Fenprostalene.

A tolerance for marker residue of fenprostalene in cattle is not needed. The safe concentrations for the total residues of fenprostalene in the uncooked edible tissues of cattle are 10 parts per billion in muscle, 20 parts per billion in liver, 30 parts per billion in kidney, 40 parts per billion in fat, and 100 parts per billion in the injection site. As used in this section “tolerance” refers to a concentration of a marker residue in the target tissue selected to monitor for total residues of the drug in the target animal, and “safe concentrations” refer to the concentrations of total residues considered safe in edible tissues.

[49 FR 26716, June 29, 1984]

§ 556.283 Florfenicol.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of florfenicol is 10 micrograms per kilogram of body weight per day.

(b) *Cattle*. A tolerance of 3.7 parts per million (ppm) for florfenicol amine (marker residue) in liver (target tissue) is established. A tolerance of 0.3 ppm for florfenicol amine in cattle muscle is established.

[63 FR 41191, Aug. 3, 1998]

§ 556.286 Flunixin meglumine.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of flunixin is 0.72 micrograms per kilogram of body weight per day.

(b) *Tolerances*. For residues of parent flunixin free acid of 0.125 part per million (ppm) in cattle liver (target tissue) and 0.025 ppm in cattle muscle are established.

[63 FR 38750, July 20, 1998]

§ 556.290 Furazolidone.

A tolerance of zero is established for residues of furazolidone in the uncooked edible tissues of swine.

§ 556.300 Gentamicin sulfate.

(a) A tolerance of 0.1 part per million is established for negligible residues of gentamicin sulfate in the uncooked edible tissues of chickens and turkeys.

(b) Tolerances are established for total residues of gentamicin in edible tissues of swine as follows: 0.1 part per million in muscle, 0.3 part per million in liver, and 0.4 part per million in fat and kidney. A microbiological determinative procedure and an HPLC confirmatory procedure for gentamicin have been developed to assay gentamicin in kidney at 0.4 ppm. Since residues of gentamicin as the parent compound and total residues are equal, the marker (parent drug) residue concentration of 0.4 ppm in kidney corresponds to 0.4 ppm of total residue.

[48 FR 791, Jan. 7, 1983, as amended at 61 FR 24441, May 15, 1996]

§ 556.308 Halofuginone hydrobromide.

The marker residue selected to monitor for total residues of halofuginone hydrobromide in broilers and turkeys is parent halofuginone hydrobromide and the target tissue selected is liver. A tolerance is established in broilers of 0.16 part per million and in turkeys of 0.13 part per million for parent halofuginone hydrobromide in liver. These marker residue concentrations in liver correspond to total residue concentrations of 0.3 part per million in liver. The safe concentrations for total residues of halofuginone hydrobromide in the uncooked edible tissues of broilers and turkeys are 0.1 part per million in muscle, 0.3 part per million in liver, and 0.2 part per million in skin with adhering fat. As used in this section, “tolerance” refers to a concentration of a marker residue in the target tissue selected to monitor for total residues of the drug in the target animal, and “safe concentrations” refers to the concentrations of total residues considered safe in edible tissues.

[54 FR 28052, July 5, 1989, as amended at 56 FR 8711, Mar. 1, 1991; 57 FR 21209, May 19, 1992]

§ 556.310 Haloxon.

A tolerance of 0.1 part per million is established for negligible residues of